



YamayBio

One-Step Stain Free PAGE Gel Rapid Preparation Kit

QUICK START GUIDE

Research use only.
Not for diagnostic procedures.

Contents and storage

| Product | Cat. No. | Quantity | Component | Cat. No. | Volume |
|--|----------|------------|----------------------------|----------|--------|
| One-Step Stain Free PAGE Gel Rapid Preparation Kit (6%) | RF1521 | 100 Assays | Stacking Buffer A | RF1501 | 80 mL |
| | | | Stacking Buffer B | RF1502 | 80 mL |
| | | | Resolving Buffer A (6%) | RF1503 | 250 mL |
| | | | Resolving Buffer B (6%) | RF1508 | 250 mL |
| | | | Enhanced Catalyst | RF1500 | 8 mL |
| One-Step Stain Free PAGE Gel Rapid Preparation Kit (7.5%) | RF1522 | 100 Assays | Stacking Buffer A | RF1501 | 80 mL |
| | | | Stacking Buffer B | RF1502 | 80 mL |
| | | | Resolving Buffer A (7.5%) | RF1504 | 250 mL |
| | | | Resolving Buffer B (7.5%) | RF1509 | 250 mL |
| | | | Enhanced Catalyst | RF1500 | 8 mL |
| One-Step Stain Free PAGE Gel Rapid Preparation Kit (10%) | RF1523 | 100 Assays | Stacking Buffer A | RF1501 | 80 mL |
| | | | Stacking Buffer B | RF1502 | 80 mL |
| | | | Resolving Buffer A (10%) | RF1505 | 250 mL |
| | | | Resolving Buffer B (10%) | RF1510 | 250 mL |
| | | | Enhanced Catalyst | RF1500 | 8 mL |
| One-Step Stain Free PAGE Gel Rapid Preparation Kit (12.5%) | RF1524 | 100 Assays | Stacking Buffer A | RF1501 | 80 mL |
| | | | Stacking Buffer B | RF1502 | 80 mL |
| | | | Resolving Buffer A (12.5%) | RF1506 | 250 mL |
| | | | Resolving Buffer B (12.5%) | RF1511 | 250 mL |
| | | | Enhanced Catalyst | RF1500 | 8 mL |
| One-Step Stain Free PAGE Gel Rapid Preparation Kit (15%) | RF1525 | 100 Assays | Stacking Buffer A | RF1501 | 80 mL |
| | | | Stacking Buffer B | RF1502 | 80 mL |
| | | | Resolving Buffer A (15%) | RF1507 | 250 mL |
| | | | Resolving Buffer B (15%) | RF1512 | 250 mL |
| | | | Enhanced Catalyst | RF1500 | 8 mL |

Storage: The enhanced catalyst should be stored at -20°C; other components are stored at 4°C for 12 months. Product shipped at ambient temperature. Once opened, the enhanced catalyst can be stored at 4°C for three months.

Introduction

This kit simplifies polyacrylamide gel preparation for standard Tris-glycine electrophoresis. These solutions are formulated to allow for a quick casting method with **no polymerization wait time** between pouring the resolving and stacking gels. The stacking gel is color-coded (red, blue, or green) for easy sample loading.

Instead of TEMED, an odorless catalyst ensures rapid polymerization (15-30 minutes at room temperature). This versatile kit produces gels suitable for both denaturing and native PAGE, with **protein bands visualized** directly under UV light (302 nm).

Quick Cast Protocol

Preparation of resolving solutions and stacking solutions for gel casting

| For 1 mm thick mini-gel | | | | | | |
|-------------------------|--------------------|--------------------|----------|-------------------|-------------------|----------|
| Gel Percentage | Resolving Buffer A | Resolving Buffer B | Catalyst | Stacking Buffer A | Stacking Buffer B | Catalyst |
| Each | 2.3 mL | 2.3 mL | 50µL | 0.75 mL | 0.75 mL | 15µL |

| For 0.75 mm thick mini-gel | | | | | | |
|----------------------------|--------------------|--------------------|----------|-------------------|-------------------|----------|
| Gel Percentage | Resolving Buffer A | Resolving Buffer B | Catalyst | Stacking Buffer A | Stacking Buffer B | Catalyst |
| Each | 1.75 mL | 1.75 mL | 35µL | 0.5 mL | 0.5 mL | 10µL |

| For 1.5 mm thick mini-gel | | | | | | |
|---------------------------|--------------------|--------------------|----------|-------------------|-------------------|----------|
| Gel Percentage | Resolving Buffer A | Resolving Buffer B | Catalyst | Stacking Buffer A | Stacking Buffer B | Catalyst |
| Each | 3.45 mL | 3.45 mL | 70µL | 1.00 mL | 1.00 mL | 20µL |

*Volumes listed are sufficient for casting one 7.4 x 8.2 cm mini-gel and can be multiplied by N (desired number of gels) to cast multiple gels at once.

Instructions:

1. Prepare the resolving gel with the desired acrylamide percentage by pipetting **equal volumes** of Resolving Buffer A and Resolving Buffer B into a clean conical tube.
2. Prepare the stacking gel by pipetting **equal volumes** of Stacking Buffer A and Stacking Buffer B into a clean conical tube.
3. Add the required volume of Enhanced Catalyst into the resolving tube. Gently mix reagents, avoiding the introduction of air bubbles into the gel mixture. Using a pipette, fill each cassette to 0.5-1 cm below the comb teeth.
4. Add the required volume of Enhanced Catalyst into the stacking tube. Gently mix reagents, avoiding the introduction of air bubbles into the gel mixture.
5. Position the pipette at the middle of the cassette and gently add the stacking gel, filling to the top of the short plate. A dip may occur where pipetting takes place but will level out.
6. Quickly and carefully insert the comb and avoid air bubble entrapment below the teeth.
7. Allow gels to polymerize for 15 minutes.
8. Gels can be used immediately or wrapped in DI water-soaked paper towels and stored in an airtight container at 4°C for up to 5 days.
9. Remove the comb before loading samples. The recommended electrophoresis conditions are **150V for 60 minutes** or **200V for 45 minutes**.
10. Following electrophoresis, carefully remove the gel cassette and glass slides. Protein bands can be visualized under **UV light** (302 nm).
11. After transferring proteins to an NC or PVDF membrane, expose it again to UV light to confirm protein transfer.
12. Please note that the excitation of fluorescent dyes may take approximately 1-5 minutes. Please be patient until the protein bands become clearly visible.
13. For optimal visualization of fluorescent protein bands on the membrane, pre-exposure to UV light may be necessary before protein transfer. This pre-excitation step can enhance fluorescence signal intensity.

Note:

1. Gel polymerization time may vary depending on temperature. Higher temperatures generally accelerate polymerization, while lower temperatures may prolong the process. In colder environments, consider increasing the Catalyst volume by up to twofold.
2. If the kit has been stored at 4°C, allow it to reach room temperature before use to minimize the formation of air bubbles during gel casting.



YamayBio